

Credit Valley Hospital 2200 Eglinton Avenue West Mississauga, Ontario L5M 2N1

DIAGNOSTIC IMAGING CONSULTATION REQUEST

Date:								
For Appointments call: (905) 813-2731 or	Fax to:	(905	5) 813-4418					
☐ Mammography ☐ Bone Density ☐ Ultrasound ☐ Nuclear Medicine ☐ G.I. Series ☐ C.T. CT-Redire ☐ 1 Trillin ☐ 1 HHS ☐ 1 HHS REGIONS TO EXAMINE ☐ 1 Any	ct to: um Heal Oakville Milton	Spo	ecial Procedur ntre	res	*PLEASE ARRIVE ON REQUISITION FOR YOU OTHERWISE WE MAY YOUR STUDY.	OUR APP	OINTMENT.	
1								
2				Refer	ring Physician:			
				Fax #	:		_	
3								
PROBLEM FOR INVESTIGATION (PLEA HISTORY / FINDINGS	SE PRI	NT C	LEARLY)					
THISTORY / TINDHNGS							-	
							_	
						==	_	
							_	
CC:	- 1 h -	4 7 G		PHYSIC	IAN'S SIGNATURE		_	
YOUR APPOINTMENT HAS BEEN SCHEDUL	ED FOR:	DAY	/	DATE	: TIME:_			
CONTRAST INJECTION CHECKLIST (To Be Completed By Imaging Personnel)								
	Yes	No				Yes	No	
Previous Contrast Examination Date:					Sickle Cell Disease Polycythemia			
					Pheochromocytoma	📋		
Previous Contrast Reaction Date:			5. Patient has	s read i	nformation in contrast			
	_		material br	rochure		🗖		
3. Allergy History: Asthma Hay Fever Drugs			6. Contrast U Amount:			. 🗖	ū	
Other			7. Injected By	V:	Ti	me:		
4. Other Risk Factors: Heart Disease								
Hypertension Diabetes	Ē							
Myeloma Renal Disease		100	9. Study Sup	ervised	By:		_MD	



PATIENT PREPARATION FOR DIAGNOSTIC IMAGING PROCEDURES

(BRACKETED TIMES = APPROXIMATE DURATION OF PROCEDURES)
For Paediatric and other examinations NOT covered below, call (905) 813-4517

NUICI EAD MEDICINE						
NUCLEAR MEDICINE .						
□ Bone Scan (4 Hours) No preparation required. □ Biliary Scan (1-2 Hours) DO NOT eat or drink 4 hours prior to examination □ Schillings Test (2 Day Test) DO NOT eat or drink 4 hours prior to the examination. Check with your doctor about stopping Vit. B12 prior to your test.	 ☐ Thyroid or Parathyroid Scans (2 Day Test) DO NOTeat or drink 4 hours prior to the examination. No x-ray contrast media ("dye") within 4 weeks of your scan; if on thyroid medication please check with your doctor ☐ Renal (Kidney) scan for High Blood Pressure Omit breakfast. Check with your doctor if you are taking blood pressure medication ☐ Renal (Kidney) scan: Routine/Diuretic Drink 4 glasses of water during the 1 hour before your test. Urinate as needed. 					
GI	СТ					
☐ Esophagus, Stomach & Duodenum G.I. Series (1 Hour) DO NOT eat or drink after 10:00 p.m. the night before before the examination.	☐ HEAD DO NOT eat or drink for 2 hours prior ☐ ABDOMINAL DO NOT eat or drink for 4 hours prior					
 ☐ <u>Small Bowel Follow Through</u> (1-3 Hours) DO NOT eat or drink after 10:00 p.m. the night before before the examination. 	MAMMOGRAPHY (30 Minutes) ☐ DO NOT use deodorant or talcum powder. Wear a 2 piece outfit.					
☐ Barium Enema (2 Hours)	GU					
Preparation pamphlet to be given by Physician's Office or picked up at hospita	☐ Intravenous Pyelogram I.V.P. (1 Hour) Clear fluid breakfast if morning appointment. Regular breakfast and clear fluid lunch if afternoon appointment.					
ULTRASOUND						
 □ Abdomen (Gall Bladder, Liver, Pancreas, Spleen, Kidney and Aorta) □ DO NOT eat or drink anything for 12 hours before. your appointment. Avoid gas producing foods such as 						
beans and carbonated beverages for 24 hours before the examination. There will be less gas if you refrain from smoking and chewing gum for 12 hours. Patients on medication may take it with a small amount of water. Insulin dependent diabetics should take their insulin with dry toast and juice.	You must administer a fleet emema to yourself 1 hour before the examination. This may be purchased at any drug store.					
Chatatrical and Palvia	NOTE:					
Obstetrical and Pelvis Your bladder must be full. Therefore 2 hours before the examination empty your bladder, then drink FOUR LARGE GLASSES of water / or some other CLEAR liquid. You must have completed drinking 1 hour before your examination.	 There is no known harmful effect to adult, child or fetus. No X-rays are used. Examination time usually varies from 30 to 60 minutes. The examination is usually not painful. 					